arXiv:1208.4433v1 [hep-th] 22 Aug 2012

DBI Galileon inflation in background SUGRA

Sayantan Choudhury[*] and Supratik Pal[†]
Physics and Applied Mathematics Unit,
Indian Statistical Institute,
203 B.T. Road, Kolkata 700 108, India

(Dated: August 23, 2012)

We introduce a model of *potential driven DBI Galileon inflation* in background $\mathcal{N}$=1, $\mathcal{D}$=4 SUGRA. Starting from D4-$\bar{D}$4 brane-antibrane in the bulk $\mathcal{N}$=2, $\mathcal{D}$=5 SUGRA including quadratic Gauss-Bonnet corrections, we derive an effective $\mathcal{N}$=1, $\mathcal{D}$=4 SUGRA by dimensional reduction, that results in a Coleman-Weinberg type Galileon potential. We employ this potential in modeling inflation and in subsequent study of primordial quantum fluctuations for scalar and tensor modes. Further, we estimate the major observable parameters in both *de Sitter (DS)* and *beyond de Sitter (BDS)* limits and confront them with recent observational data from WMAP7 by using the publicly available code CAMB.

Contents

I. INTRODUCTION 1
II. The Background Action 2
III. Generation of D3 DBI Galileon 5
  A. **The Einstein-Hilbert Action** 5
  B. **The Gauss-Bonnet Action** 5
  C. **The D3 Brane Action** 6
  D. **The $\mathcal{N}$=1, $\mathcal{D}$=4 Supergravity Action** 6
IV. Modeling Inflation from DBI Galileon 7
V. Quantum Fluctuations and Observable Parameters 10
VI. Parameter Estimation and Confrontation with WMAP7 13
VII. Summary and Outlook 14
Acknowledgments 15
Appendix 15
References 18

I. INTRODUCTION

In the very recent days a good number of theoretical physicists have devoted their attention to the development of consistent modified gravity theories which play analogous role as dark energy or the cosmological constant [1], [2]. In higher-dimensional setups as in the case of *DGP model* [3],[4] where self-acceleration is sourced by a scalar field (the zero helicity mode of the 5D graviton), these types of Infra-Red (IR) modification of gravity [5], [6] play a crucial role. Moreover, the *DGP model* replicates the general relativistic features due to non-linear interactions via the well known *Vainshtein mechanism* [7]- [9]. Despite its profound success it has got some serious limitations

---

[*] Electronic address: sayanphysicsisi@gmail.com
[†] Electronic address: supratik@isical.ac.in

[8], [10], which are resolved by introducing a dynamical field, *aka*, Galileon [11]-[13] arising on the brane from the bulk in the DGP setup. The cosmological consequences of the Galileon models have been studied to some extent in [14]-[16]. Very recently, a natural extension to the scenario has been brought forward by tagging Galileon with the good old DBI model [17], [18], resulting in so-called "DBI Galileon" [19]-[21], that has reflected a rich structure from four dimensional cosmological point of view. However, in most of the physical situations, this type of *effective* gravity theories are plagued with additional degrees of freedom which often results in unwanted debris like ghosts, Laplacian instabilities etc [22], [23]. In the present paper we introduce a single scalar field model described by the D3 DBI Galileon originated from D4-$\bar{D}4$ brane anti-brane setup. This prevents the framework from having extra degrees of freedom as well as *Ostrogradski instabilities* [24], resulting in a higher-order derivative scalar field theory free from any such unwanted instabilities. Nevertheless, a consistent field theoretic derivation of the effective potential commonly used in the context of DBI Galileon cosmology has not been brought forth so far. On top of that, it is imperative to point out that the SUGRA origin of D3 DBI Galileon is yet to be addressed. In this article we plan to address both of these issues explicitly by deriving the inflaton potential from our proposed framework of DBI Galileon. It turns out that the derived action includes, in certain limits, the decoupling limit of DGP model as well as some consistent theories of massive gravity; and it also includes the "K-mouflage" [25], [26] and also "G/KGB" inflation [16], [27]-[29]. Moreover, in general appearance of non-vanishing superstringy frame functions in the 4D action expedites breakdown of shift symmetry. Without shift symmetry, it may happen that the theory is unstable against large renormalization. The background action chosen in our model preserves shift symmetry of the single scalar field which gives it a firm footing from phenomenological point of view as well.

The plan of the paper is as follows. First we propose a fairly general framework by taking the full DBI action in D4 brane in the background of bulk $\mathcal{N}$=2, $\mathcal{D}$=5 supergravity [30]-[32] including the quadratic modification in Einstein's Hilbert action via Gauss-Bonnet correction in the bulk. Hence, using dimensional reduction technique, we derive the effective action for DBI Galileon in D3 brane in the background of $\mathcal{N}$=1, $\mathcal{D}$=4 supergravity induced by the quadratic correction in the geometry sector and study cosmological inflationary scenario therefrom. We next engage ourselves in studying quantum fluctuations, by employing second order perturbative action for scalar and tensor modes in *de Sitter (DS)* and *beyond de Sitter (BDS)* limits, and hence calculate the primordial power spectrum of the scalar and tensor modes, their running and other observable parameters. We further confront our model with observations by using the publicly available code CAMB [33], and find them to fit well with latest observational data from WMAP7[34] and expected to fit fair well with upcoming data from PLANCK[35].

II. THE BACKGROUND ACTION

Let us demonstrate briefly the construction of DBI Galileon starting from $\mathcal{N}$=2,$\mathcal{D}$=5 SUGRA along with Gauss Bonnet correction in D4 brane set up. The full five dimensional model is described by the following action

$$S^{(5)}_{Total} = S^{(5)}_{EH} + S^{(5)}_{GB} + S^{(5)}_{D4\ brane} + S^{(5)}_{BulkSugra} \tag{2.1}$$

where

$$S^{(5)}_{EH} = \frac{1}{2\kappa_5^2}\int d^5x\sqrt{-g^{(5)}}\left[R_{(5)} - 2\Lambda_5\right], \tag{2.2}$$

$$S^{(5)}_{GB} = \frac{\alpha_{(5)}}{2\kappa_5^2}\int d^5x\sqrt{-g^{(5)}}\left[R^{ABCD(5)}R^{(5)}_{ABCD} - 4R^{AB(5)}R^{(5)}_{AB} + R^2_{(5)}\right]. \tag{2.3}$$

The D4 brane action decomposed into two parts as

$$S^{(5)}_{D4\ brane} = S^{(5)}_{DBI} + S^{(5)}_{WZ}, \tag{2.4}$$

where the *DBI* action and the *Wess-Zumino* action are given by respectively

$$S^{(5)}_{DBI} = -\frac{T_4}{2}\int d^5x\, \exp(-\Phi)\sqrt{-\left(\gamma^{(5)} + B^{(5)} + 2\pi\alpha' F^{(5)}\right)}, \tag{2.5}$$

$$\begin{aligned} S^{(5)}_{WZ} &= -\frac{T_4}{2}\int \sum_{n=0,2,4} \hat{C}_n \wedge \exp\left(\hat{B}_2 + 2\pi\alpha^{'} F_2\right)|_{4\ form} \\ &= \frac{1}{2}\int d^5x\sqrt{-g^{(5)}}\left\{\epsilon^{ABCD}\left[\partial_A\Phi^I\partial_B\Phi^J\left(\frac{C_{IJ}B_{KL}}{4T_4}\partial_C\Phi^K\partial_D\Phi^L + \frac{\pi\alpha^{'}C_{IJ}F_{CD}}{2}\right.\right.\right. \\ &\left.\left.\left. + \frac{C_0}{8T_4}B_{IJ}B_{KL}\partial_C\Phi^K\partial_D\Phi^L + \frac{\pi\alpha^{'}C_0}{2}B_{IJ}F_{CD}\right) + 2\pi^2\alpha^{'2}T_4C_0F_{AB}F_{CD} - T_4\nu(\Phi)\right]\right\} \end{aligned} \tag{2.6}$$

where $T_{(4)}$ is the D4 brane tension, $\alpha^{'}$ is the Regge Slope, $\exp(-\Phi)$ is the closed string dilaton and C_0 is the Axion. The gauge invariant combination of rank 2 field strength tensor, appearing in D4 brane, is $\mathcal{F}_{AB} = B_{AB} + 2\pi\alpha^{'}F_{AB}$ and $\{F_2, B_2\}$ represents 2-form $U(1)$ gauge fields. In eqn(2.1) $\mathcal{N} = 2, \mathcal{D} = 5$ bulk supergravity action can be written as [30]

$$\begin{aligned} S^{(5)}_{Bulk\ Sugra} = \tfrac{1}{2}\int d^5x\sqrt{-g_{(5)}}e_{(5)}\left[-\tfrac{M_5^3R^{(5)}}{2} + \tfrac{i}{2}\bar{\Psi}_{i\tilde{m}}\Gamma^{\tilde{m}\tilde{n}\tilde{q}}\nabla_{\tilde{n}}\Psi^i_{\tilde{q}} - S_{IJ}F^I_{\tilde{m}\tilde{n}}F^{I\tilde{m}\tilde{n}} - \tfrac{1}{2}g_{\alpha\beta}(D_{\tilde{m}}\phi^\mu)(D^{\tilde{m}}\phi^\nu)\right] \\ + \mathrm{Fermionic} + \mathrm{Chern-Simons} + \mathrm{Pauli\ \ mass}, \end{aligned} \tag{2.7}$$

In this context the 5-dimensional coordinates $X^A = (x^\alpha, y)$, where y parameterizes the extra dimension compactified on the closed interval $[-\pi R, +\pi R]$ and Z_2 symmetry is imposed. We consider 5D *Yang Mills* SUGRA model which is described by the field content $\{e^{\tilde{m}}_{\tilde{\mu}}, \Psi^i_{\tilde{\mu}}, A^I_{\tilde{\mu}}, \lambda^{ia}, \Phi^x\}$ where $\tilde{\mu} = (\mu, 5)$ are curved and $\tilde{m} = \left(m, \dot{5}\right)$ are flat 5D indices with μ, m their corresponding 4D indices. The remaining indices are $I = 0, 1, ..., n$, $a = 1, 2, ..., n$ and $x = 1, 2, ..., n$. The SUGRA multiplet consists of the f$\ddot{u}$nfbien $e^{\tilde{m}}_{\tilde{\mu}}$, two graviphoton $A^0_{\tilde{\mu}}$ and two gravitini $\Psi^i_{\tilde{\mu}}$, where $i = 1, 2$ is the simplectic $SU(2)_R$ index. Moreover, there exists n vector multiplets, counting the *Yang Mills* fields $(A^a_{\tilde{\mu}})$. The spinor and the scalar fields included in the vector multiplets are collectively denoted by λ^{ia}, Φ^x respectively. The indices a and x are flat and curved indices respectively of the 5D manifold $\mathcal{M}$. It is important to mention here that the Chern-Simons terms can be gauged away assuming cubic constraints and Z_2 symmetry. Now we consider full particle spectrum , the Z_2 even fields $\{e^m_\mu, e^{\dot{5}}_5, \Psi^1_\mu, \Psi^2_5, A^0_\mu, \lambda^{1a}\}$ and the Z_2 odd fields $\{e^{\dot{5}}_\mu, e^m_5, \Psi^2_\mu, \Psi^1_5, A^0_\mu, A^a_5, \lambda^{2a}, \Phi^x\}$ propagates in the bulk. For computational purpose it is useful to define the five dimensional generating function(G) of supergravity in this setup as

$$G = -3\ln\left(\frac{T + T^\dagger}{\sqrt{2}}\right) + K(\Phi, \Phi^\dagger), \tag{2.8}$$

where the supergravity K$\ddot{a}$hler moduli fields are given by $T = \frac{(e^{\dot{5}} - i\sqrt{\frac{2}{3}}A^0_5)}{\sqrt{2}}$ which is assumed to be stabilized under first approximation and $K(\Phi, \Phi^\dagger)$ represents generalized K$\ddot{a}$hler function.

Including the kinetic term of the five dimensional field Φ and rearranging into a perfect square, the 5D bulk supergravity action can be expressed as

$$S \supset \frac{1}{2}\int d^4x\int_{-\pi R}^{+\pi R} dy\sqrt{-g_5}e_{(4)}e^5_{\dot{5}}\left[g^{\alpha\beta}G^N_M(\partial_\alpha\Phi^M)^\dagger(\partial_\beta\Phi_N) + \frac{1}{g_{55}}\left(\partial_5\phi - \sqrt{V^{(5)}_{bulk}(G))}\right)^2\right], \tag{2.9}$$

where the 5D potential

$$V^{(5)}_{bulk}(G) = \exp\left(\frac{G}{M^2}\right)\left[\left(\frac{\partial W}{\partial\phi_M} + \frac{\partial G}{\partial\phi_M}\frac{W}{M^2}\right)^\dagger(G^N_M)^{-1}\left(\frac{\partial W}{\partial\phi^N} + \frac{\partial G}{\partial\phi^N}\frac{W}{M^2}\right) - 3\frac{|W|^2}{M^2}\right]. \tag{2.10}$$

The field equations in presence of Gauss-Bonnet term can be expressed as

$$G^{(5)}_{AB} + \alpha_{(5)}H^{(5)}_{AB} = 8\pi G_{(5)}T^{(5)}_{AB} - \Lambda_{(5)}g^{(5)}_{AB}, \tag{2.11}$$

where the covariantly conserved Gauss-Bonnet tensor

$$H^{(5)}_{AB} = 2R^{(5)}_{ACDE}R^{CDE(5)}_B - 4R^{(5)}_{ACBD}R^{CD(5)} - 4R^{(5)}_{AC}R^{C(5)}_B + 2R^{(5)}R^{(5)}_{AB} - \frac{1}{2}g^{(5)}_{AB}\left(R^{ABCD(5)}R^{(5)}_{ABCD} - 4R^{AB(5)}R^{(5)}_{AB} + R^2_{(5)}\right) \tag{2.12}$$

which acts as a source term. It is useful to introduce the 5D metric in conformal form

$$ds^2_{4+1} = g_{AB}dX^A dX^B = \frac{1}{\sqrt{h(y)}}ds^2_4 + \sqrt{h(y)}\tilde{G}(y)dy^2 = \exp(2A(y))\left(ds^2_4 + R^2\beta^2 dy^2\right), \tag{2.13}$$

with warp factor

$$\exp(2A(y)) = \frac{1}{\sqrt{h(y)}} = \frac{R^2}{b_0^2\beta^2}\tilde{G}(y) = \frac{b_0^2}{R^2\left(\exp(\beta y) + \frac{\Lambda_{(5)}b_0^4}{24R^2}\exp(-\beta y)\right)} \tag{2.14}$$

and $ds^2_4 = g_{\alpha\beta}dx^\alpha dx^\beta$ is FLRW counterpart. In order to write down explicitly the expression for D4 brane action, the induced metric can be shown as

$$\gamma_{CD} = \frac{1}{\sqrt{h(y)}}\left(g_{AB} + h(y)\tilde{G}_{AB}\partial_C\Phi^A\partial_D\Phi^B\right). \tag{2.15}$$

The 5D energy momentum tensor for the set up reads

$$\begin{aligned}T^{total(5)}_{\alpha\beta} = T^{bulk(5)}_{\alpha\beta} + T^{D4brane(5)}_{\alpha\beta} = {} & G^N_M(\partial_\alpha\Phi^M)^\dagger(\partial_\beta\Phi_N) - g_{\alpha\beta}\left[g^{\rho\sigma}(\partial_\rho\Phi^M)^\dagger(\partial_\sigma\Phi_N)G^N_M + g^{55}\left(\partial_5\Phi - \sqrt{V^{(5)}_{bulk}(G)}\right)^2\right] \\ & + \left[K_{MN;X}\partial_\alpha\Phi^M\partial_\beta\Phi^N + Kg_{\alpha\beta} - 2\nabla_{(\alpha}G(\Phi,X)\nabla_{\beta)}\Phi + g_{\alpha\beta}\partial_\lambda G(\Phi,X)\partial^\lambda\Phi \right.\\ & \left. - G_{MN;X}(\Phi,X)\Box\Phi\partial_\alpha\Phi^M\partial_\beta\Phi^N\right],\end{aligned} \tag{2.16}$$

$$\begin{aligned}T^{total(5)}_{55} = T^{bulk(5)}_{55} + T^{D4brane(5)}_{55} = {} & \tfrac{1}{2}\left(\partial_5\Phi - \sqrt{V^{(5)}_{bulk}(G)}\right)^2 - \tfrac{1}{2}g_{55}g^{\rho\sigma}G^N_M(\partial_\rho\Phi^M)^\dagger(\partial_\sigma\Phi_N) \\ & + \left[K_{MN;X}\partial_5\Phi^M\partial_5\Phi^N + Kg_{55} - 2\nabla_{(5}G(\Phi,X)\nabla_{5)}\Phi + g_{55}\partial_\lambda G(\Phi,X)\partial^\lambda\Phi \right.\\ & \left. - G_{MN;X}(\Phi,X)\Box\Phi\partial_5\Phi^M\partial_5\Phi^N\right].\end{aligned} \tag{2.17}$$

On the other hand, the *Klein-Gordon* equation of motion in 5D can be expressed as

$$\begin{aligned}&\partial_5\left[\tfrac{e^5_{\dot{5}}\sqrt{g_5}}{g_{55}}\left(\partial_5\Phi - \sqrt{V^{(5)}_{bulk}(G)}\right)\right] + \textstyle\sum_N e^5_{\dot{5}}\left\{\partial_\beta\left[\sqrt{g_5}g^{\alpha\beta}G^N_M(\partial_\alpha\Phi^M)\right] - \tfrac{\sqrt{g_5}}{g_{55}}\partial_N\left(\sqrt{V^{(5)}_{bulk}(G)}\right)\left(\partial_5\Phi - \sqrt{V^{(5)}_{bulk}(G)}\right)\right\} \\ &K_X(\Phi,X)\Box^{(5)}\Phi - K_{XX}(\Phi,X)\left(\nabla_A\nabla_B\Phi\right)\left(\nabla^A\Phi\nabla^B\Phi\right) - 2K_{\Phi X}(\Phi,X)X + K_\Phi(\Phi,X) - 2\left(G_\Phi(\Phi,X) - G_{\Phi X}(\Phi,X)X\right) \\ &+G_X(\Phi,X)\left[\left(\nabla_A\nabla_B\Phi\right)\left(\nabla^A\nabla^B\Phi\right) - \left(\Box^{(5)}\Phi\right)^2 + R^{(5)}_{AB}\nabla^A\Phi\nabla^B\Phi\right] + 2G_{\Phi X}(\Phi,X)\left(\nabla_A\nabla_B\Phi\right)\left(\nabla^A\nabla^B\Phi\right) + 2G_{\Phi\Phi}(\Phi,X)X \\ &-G_{XX}\left(\nabla^A\nabla^B\Phi - g^{AB}\Box^{(5)}\Phi\right)\left(\nabla_A\nabla^C\Phi\right)\left(\nabla_B\Phi\right)\left(\nabla_C\Phi\right) = 0.\end{aligned} \tag{2.18}$$

Now using the scaling relations $\Phi^A = \sqrt{T_{(4)}}\tilde{\Phi}^A$, $G_{AB} = \exp(-\Phi)\tilde{g}_{AB}$ and $b_{AB} = \frac{\sqrt{h(y)}}{T_{(4)}}B_{AB}$ the 5D action for D4 brane can be expressed in more convenient form as

$$S^{(5)}_{D4\ brane} = \int d^5x\sqrt{-g^{(5)}}\left[K(\Phi,X) - G(\Phi,X)\Box^{(5)}\Phi\right], \tag{2.19}$$

where $K(\Phi,X) = -\frac{1}{2f(\Phi)}\left(\sqrt{\mathcal{D}} - 1\right) - \frac{V^{(5)}_{brane}(\Phi)}{2}$, the determinant $\mathcal{D} = 1 - 2f(\Phi)G_{AB}X^{AB} + 4f^2(\Phi)X^{[A}_A X^{B]}_B - 8f^3(\Phi)X^{[A}_A X^B_B X^{C]}_C + 16f^4(\Phi)X^{[A}_A X^B_B X^C_C X^{D]}_D$, kinetic term $X^B_D = -\frac{1}{2}G_{DA}\partial^C\Phi^A\partial_C\Phi^B$ and the 5D D'Alembertian Operator $\Box^{(5)} = \frac{1}{\sqrt{-g^{(5)}}}\partial_A\left(\sqrt{-g^{(5)}}g^{AC}\partial_C\right)$. Here we use the fact that no spatial direction along which the scalar fields are only time dependent lead to $B^\mu_\nu = 0$ and $F_{\mu\nu} = 0$ in the background. Consequently *Maxwell's field equations* are unaffected in 4D after dimensional reduction. In this context the 5D D4 brane potential is given by

$$V^{(5)}_{brane}(\Phi) = T_{(4)}\nu(\Phi) + \frac{1}{f(\Phi)}, \tag{2.20}$$

where *Klebanov Strassler* warped geometry motivated frame function $f(\Phi) = \frac{\exp(\Phi)h(y)}{T_{(4)}} \simeq \frac{1}{(f_0 + f_2\Phi^2 + f_4\Phi^4)}$ (using Z_2) is originated from higher dimensional superstring theory, *Coulomb* frame function $\nu(\Phi) = \left(\nu_0 + \frac{\nu_4}{\Phi^4}\right)$ is originated from Coulomb interaction between D4-$\bar{D}$4 brane and the implicit brane function $G(\Phi,X) = \frac{g(\Phi)}{2(1-2f(\Phi)X)}$ with $g(\Phi) = g_0 + g_2\Phi^2$.

III. GENERATION OF D3 DBI GALILEON

In this section we employ dimensional reduction technique to derive a $\mathcal{N}$=1, $\mathcal{D}$=4 SUGRA and the inflaton potential therefrom that results in DBI Galileon on the D3 brane. For convenience we deal with different contributions to the action (2.1) separately.

A. The Einstein-Hilbert Action

After integrating out the contribution from the five dimension, the Einstein Hilbert action in four dimension can be written as

$$\begin{aligned} S^{(4)}_{EH} &= \frac{1}{2}\int d^4x\sqrt{-g_{(4)}}\int_{-\pi R}^{+\pi R} dy\beta M_5^3 R\exp(3A(y))\left[R_{(4)} - \frac{12}{\beta^2 R^2}\left(\frac{dA(y)}{dy}\right)^2 - \frac{8}{\beta^2 R^2}\left(\frac{d^2A(y)}{dy^2}\right) - 2\Lambda_5\exp(2A(y))\right] \\ &= \frac{1}{2\kappa_4^2}\int d^4x\sqrt{-g_{(4)}}\left[R_{(4)} - \frac{3M_5^3\beta b_0^6}{M_{PL}^2 R^5}\mathcal{I}(1)\right], \end{aligned} \tag{3.1}$$

where the explicit expression for $\mathcal{I}(1)$ is mentioned in appendix. It is important to mention here that the 5D Planck mass and 4D Planck mass are related through

$$\begin{aligned} M^2_{PL} &= M^3_{(5)}\beta R\int_{-\pi R}^{+\pi R} dy\exp(3A(y)) \\ &= \frac{M^3_{(5)}b_0^3}{3R^2T^{3/2}_{(4)}}\left[\exp(\beta\pi R)\left\{\frac{3\sqrt{T_{(4)}}}{\sqrt{\exp(\beta\pi R)+T_{(4)}\exp(-\beta\pi R)}} - \sqrt{\frac{\exp(2\pi\beta R)+T_{(4)}}{\exp(\beta\pi R)+T_{(4)}\exp(-\beta\pi R)}}\ {}_2F_1\left[\frac{1}{2};\frac{3}{7};\frac{7}{4};-\frac{\exp(2\beta\pi R)}{T_{(4)}}\right]\right\}\right. \\ &\qquad \left. - \exp(-\beta\pi R)\left\{\frac{3\sqrt{T_{(4)}}}{\sqrt{\exp(-\beta\pi R)+T_{(4)}\exp(\beta\pi R)}} - \sqrt{\frac{\exp(-2\beta\pi R)+T_{(4)}}{\exp(-\beta\pi R)+T_{(4)}\exp(\beta\pi R)}}\ {}_2F_1\left[\frac{1}{2};\frac{3}{7};\frac{7}{4};-\frac{\exp(-2\beta\pi R)}{T_{(4)}}\right]\right\}\right]. \end{aligned} \tag{3.2}$$

B. The Gauss-Bonnet Action

The 5D and 4D *Riemann tensor*, *Ricci tensor* and *Ricci scalar* are related through the following expressions:

$$R^{(5)}_{\alpha\beta\gamma\delta} = R^{(4)}_{\alpha\beta\gamma\delta} + \frac{\exp(2A(y))}{R^2\beta^2}\left(\frac{dA(y)}{dy}\right)^2\left[g^{(4)}_{\gamma\beta}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\beta}\right] \tag{3.3}$$

$$R^{(5)}_{\alpha\beta} = R^{(4)}_{\alpha\beta} - \frac{3g^{(4)}_{\alpha\beta}}{R^2\beta^2}\left(\frac{dA(y)}{dy}\right)^2 \tag{3.4}$$

$$R_{(5)} = \exp(2A(y))\left[R_{(4)} - \frac{12}{\beta^2R^2}\left(\frac{dA(y)}{dy}\right)^2 - \frac{8}{\beta^2R^2}\left(\frac{d^2A(y)}{dy^2}\right) - 2\Lambda_5\exp(2A(y))\right] \tag{3.5}$$

Using eqn(3.3)-eqn(3.5) in eqn(2.3) we get

$$\begin{aligned} S^{(4)}_{GB} = \frac{\alpha_{(4)}}{2\kappa^2_{(4)}}\int d^4x\sqrt{-g_{(4)}}\Bigg[&\left(\mathcal{C}(1)R^{\alpha\beta\gamma\delta(4)}R^{(4)}_{\alpha\beta\gamma\delta} - 4\mathcal{I}(2)R^{\alpha\beta(4)}R^{(4)}_{\alpha\beta} + \mathcal{A}(6)R^2_{(4)}\right) \\ &+ \frac{2\mathcal{C}(2)}{R^2\beta^2}R^{(4)}_{\alpha\beta\gamma\delta}\left(g^{\gamma\beta(4)}g^{\delta\alpha(4)} - g^{\gamma\alpha(4)}g^{\delta\beta(4)}\right) + \frac{\mathcal{G}(1)}{R^4\beta^4} + \frac{\mathcal{G}(2)}{R^2\beta^2}R_{(4)}\Bigg] \end{aligned} \tag{3.6}$$

where $\mathcal{G}(1) = 24\mathcal{C}(4) - 144\mathcal{I}(4) - 64\mathcal{A}(5) + 144\mathcal{A}(7) + 64\mathcal{A}(8) + 192\mathcal{A}(11)$, $\mathcal{G}(2) = 24\mathcal{I}(2) - 24\mathcal{A}(9) - 16\mathcal{A}(10)$ and the scaling relationship between 4D and 5D *Gauss-Bonnet* coupling constant is $\alpha_{(4)} = \frac{\kappa^2_{(4)}\beta R}{\kappa^2_{(5)}}\alpha_{(5)}$. Explicit form of each of the constants appearing in eqn(3.6) is mentioned in the appendix.

C. The D3 Brane Action

To reduce the D4 brane action we employ the method of separation of variable $\Phi(X^A) = \Phi(x^\mu, y) = \tilde{\phi}(x^\mu)\chi(y)$ where $\chi(y) = \exp(\frac{2\pi i y}{R})$. Consequently the D3 brane action turns out to be

$$S^{(4)}_{D3\ Brane} = \int d^4x\sqrt{-g^{(4)}}\left[\tilde{K}(\tilde{\phi}, \tilde{X}) - \tilde{G}(\tilde{\phi}, \tilde{X})\Box^{(4)}\tilde{\phi}\right], \tag{3.7}$$

where $\tilde{K}(\tilde{\phi}, \chi) = \left\{-\frac{\tilde{D}}{\tilde{f}(\tilde{\phi})}\left[\sqrt{1 - 2Q\tilde{X}\tilde{f}} - Q_1\right] - \tilde{C}_5\tilde{G}(\tilde{\phi}, \tilde{X}) - \bar{Q}_2\tilde{D}V^{(4)}_{brane}\right\}$, $\tilde{G}(\tilde{\phi}, \tilde{X}) = \left(\frac{\tilde{g}(\tilde{\phi})k_1\tilde{C}_4}{2(1-2\tilde{f}(\tilde{\phi})\tilde{X}k_2))}\right)$, $\tilde{g}(\tilde{\phi}) = \tilde{g}_0 + \tilde{g}_2\tilde{\phi}^2$, $\tilde{D} = \frac{D}{2\kappa^2_{(4)}}$, $\tilde{C}_4 = \frac{C_4}{2\kappa^2_4}$, $\tilde{C}_5 = \frac{\bar{C}_5\beta^2R^2}{2\kappa^2_4}$, $\bar{Q}_2\tilde{D} = \beta R$. The effective *Klebanov Strassler* and *Coulomb* frame function are hereby expressed as $\tilde{f}(\tilde{\phi}) \simeq \frac{1}{(\tilde{f}_0 + \tilde{f}_2\tilde{\phi}^2 + \tilde{f}_4\tilde{\phi}^4)}$ and $\nu^{(4)}(\tilde{\phi}) = \tilde{\nu}_0 + \frac{\tilde{\nu}_4}{\tilde{\phi}^4}$ with $\tilde{\nu}_0 = \nu_0\mathcal{A}(1)$, $\tilde{\nu}_4 = \nu_4\mathcal{A}(12)$. The outcome of dimensional reduction is reflected through the constants mentioned in the appendix. The scaled D3 brane potential turns out to be

$$\tilde{V}^{(4)}_{brane} = \bar{Q}_2\tilde{D}V^{(4)}_{brane} = T_{(3)}\nu^{(4)}(\tilde{\phi}) + \frac{\beta R\mathcal{I}(2)}{\tilde{f}(\tilde{\phi})} \tag{3.8}$$

where the D3 brane tension is given by $T_{(3)} = \beta R T_{(4)}$.

D. The $\mathcal{N}$=1, $\mathcal{D}$=4 Supergravity Action

Further, imposing Z_2 symmetry to $\tilde{\phi}$ via $\Phi(0) = \Phi(\pi R) = 0$ and compactifying around a circle (S^1) $\partial_5\Phi = \sqrt{V^{(5)}_{bulk}(G)}\,(1 - \frac{1}{2\pi R})$ we get,

$$S^{(5)}_{Bulk\ Sugra} = \frac{1}{2}\int d^4x\int_{-\pi R}^{+\pi R} dy\sqrt{-g^{(5)}}\left[e_{(4)}e^5_5\left\{g^{\alpha\beta}G^n_m(\partial_\alpha\tilde{\phi}^m)^\dagger(\partial_\beta\tilde{\phi}_n) - g^{55}\frac{V^{(5)}_{bulk}(G)}{4\pi^2R^2}\right\}\right]. \tag{3.9}$$

Now using the above mentioned ansatz for method of separation of variable we get

$$S^{(4)}_{Sugra} = \frac{1}{2\kappa^2_{(4)}}\int d^4x\sqrt{-g^{(4)}}\left[M(T, T^\dagger)\mathcal{J}^\mu_\nu(\tilde{\phi}, \tilde{\phi}^\dagger)g^{\alpha\beta(4)}(\partial_\alpha\tilde{\phi}_\mu)^\dagger(\partial_\beta\tilde{\phi}^\nu) - Z(T, T^\dagger)V^{(4)}_F(\tilde{\phi})\right]. \tag{3.10}$$

where $\mathcal{J}^\mu_\nu(\tilde{\phi}, \tilde{\phi}^\dagger) = \int_{-\pi R}^{+\pi R} dy\exp(-A(y))\left(\frac{\partial^2\mathcal{K}(\tilde{\phi}\exp(\frac{2\pi i y}{R}), \tilde{\phi}^\dagger\exp(-\frac{2\pi i y}{R}))}{\partial\tilde{\phi}^\dagger_\mu\partial\tilde{\phi}^\nu}\right)$, $M(T, T^\dagger) = \frac{\sqrt{2}\beta R^2}{(T+T^\dagger)}$, $Z(T, T^\dagger) = \frac{1}{8\pi^2R^2\beta|T+T^\dagger|^2}$. Here we have used $\mathcal{W}(\tilde{\phi}, \tilde{\phi}^\dagger, T, T^\dagger) = \frac{\mathcal{W}_1(\tilde{\phi}, \tilde{\phi}^\dagger)|T+T^\dagger|^2}{4}$ for superpotential. Hence, the $\mathcal{N} = 1, \mathcal{D} = 4$ supergravity F-term potential turns out to be

$$\begin{aligned} V^{(4)}_F = \int_{-\pi R}^{+\pi R} & dy\exp(-A(y))\exp\left(\frac{\mathcal{K}(\tilde{\phi}\exp(\frac{2\pi i y}{R}), \tilde{\phi}^\dagger\exp(-\frac{2\pi i y}{R}))}{M^2}\right) \\ & \left[\left(\frac{\partial\mathcal{W}_1}{\partial\tilde{\phi}_\alpha} + \left(\frac{\partial\mathcal{K}(\tilde{\phi}\exp(\frac{2\pi i y}{R}), \tilde{\phi}^\dagger\exp(-\frac{2\pi i y}{R}))}{\partial\tilde{\phi}_\alpha}\right)\frac{\mathcal{W}_1}{M^2}\right)^\dagger\left(\frac{\partial^2\mathcal{K}(\tilde{\phi}\exp(\frac{2\pi i y}{R}), \tilde{\phi}^\dagger\exp(-\frac{2\pi i y}{R}))}{\partial\tilde{\phi}^\alpha\partial\tilde{\phi}^\dagger_\beta}\right)^{-1}\right. \\ & \left.\left(\frac{\partial\mathcal{W}_1}{\partial\tilde{\phi}^\beta} + \left(\frac{\partial\mathcal{K}(\tilde{\phi}\exp(\frac{2\pi i y}{R}), \tilde{\phi}^\dagger\exp(-\frac{2\pi i y}{R}))}{\partial\tilde{\phi}^\beta}\right)\frac{\mathcal{W}_1}{M^2}\right) - 3\frac{|\mathcal{W}_1|^2}{M^2}\right]. \end{aligned} \tag{3.11}$$

Now using the ansatz for the *Kähler* potential $\mathcal{K}(\tilde{\phi}\exp(\frac{2\pi i y}{R}), \tilde{\phi}^\dagger\exp(-\frac{2\pi i y}{R})) = \mathcal{K}_1(\tilde{\phi}, \tilde{\phi}^\dagger)\mathcal{K}_2(\exp(\frac{2\pi i y}{R}), \exp(-\frac{2\pi i y}{R}))$ with $\mathcal{K}_1(\tilde{\phi}, \tilde{\phi}^\dagger) = \mathcal{K}^{\alpha\beta}_1\tilde{\phi}_\alpha\tilde{\phi}^\dagger_\beta$ and $\mathcal{K}_2(\exp(\frac{2\pi i y}{R}), \exp(-\frac{2\pi i y}{R})) = 1$ eqn(3.11) reduces to the following form:

$$V^{(4)}_F = \mathcal{A}(13)\exp\left(\frac{\mathcal{K}^{\alpha\beta}_1\tilde{\phi}_\alpha\tilde{\phi}^\dagger_\beta}{M^2}\right)\left[\left(\frac{\partial\mathcal{W}_1}{\partial\tilde{\phi}_\alpha} + \mathcal{K}^{\alpha\beta}_1\tilde{\phi}^\dagger_\beta\frac{\mathcal{W}_1}{M^2}\right)^\dagger\mathcal{K}^\nu_{1\alpha}\left(\frac{\partial\mathcal{W}_1}{\partial\tilde{\phi}^\nu} + \mathcal{K}_{1\nu\eta}\tilde{\phi}^\eta\frac{\mathcal{W}_1}{M^2}\right) - 3\frac{|\mathcal{W}_1|^2}{M^2}\right] \tag{3.12}$$

with the general *Kähler metric* $\mathcal{K}_1^{\alpha\beta} = \frac{\partial^2 \mathcal{K}_1}{\partial\tilde{\phi}_\alpha \partial\tilde{\phi}^\dagger_\beta}$. In most of the simple situations, we are interested in the *Canonical metric* structure defined by $\mathcal{K}_1^{\alpha\beta} = \delta^{\alpha\beta}$. Consequently the $\mathcal{N}=1, \mathcal{D}=4$ SUGRA action turns out to be

$$S^{(4)}_{Can\ Sugra} = \frac{1}{2\kappa^2_{(4)}} \int d^4x \sqrt{-g^{(4)}} \left[M(T,T^\dagger) g^{\alpha\beta(4)} (\partial_\alpha \tilde{\phi}_\mu)^\dagger (\partial_\beta \tilde{\phi}^\mu) - Z(T,T^\dagger) V^{(4)}_{Can}(\tilde{\phi})\right]. \tag{3.13}$$

where the canonical F-term potential can be recast as

$$V^{(4)} = V^{(4)}_F = \mathcal{A}(13) \exp\left(\frac{\tilde{\phi}^\dagger_\alpha \tilde{\phi}^\alpha}{M^2}\right) \left[\left|\frac{\partial \mathcal{W}_1}{\partial \tilde{\phi}_\beta}\right|^2 - 3\frac{|\mathcal{W}_1|^2}{M^2}\right]. \tag{3.14}$$

To derive the expression for inflaton potential we start with a specific superpotential [36] $\mathcal{W}_1 = v\tilde{\phi} - \frac{g}{n+1}\tilde{\phi}^{n+1}$, where $\tilde{\phi}$ is the *chiral* superfield with R charge $\frac{2}{n+1}$ introduced as an inflaton with $n \geq 2$. Here $g(\sim \mathcal{O}(1))$ is the positive and real coupling constant and v is the VEV of the field $\tilde{\phi}$. In this model $U(1)_R$ symmetry is dynamically broken to a discrete $Z_{2n}R$ symmetry at the scale $v << 1$. Consequently the inflaton transforms as $\tilde{\phi} \longrightarrow \exp\left(\frac{2\alpha i}{n+1}\right)\tilde{\phi}(\exp(-i\alpha)\theta)$. This leads to the following form of the bulk contribution to the potential

$$V^{(4)}(\tilde{\phi}) = \mathcal{A}(13) \exp\left(|\tilde{\phi}|^2\right) \left[\left|\left(1+|\tilde{\phi}|^2\right)v^2 - \left(1+\frac{|\tilde{\phi}|^2}{n+1}\right) g\tilde{\phi}^n\right|^2 - 3|\tilde{\phi}|^2 \left|v^2 - \frac{g}{n+1}\tilde{\phi}^n\right|^2\right]. \tag{3.15}$$

It has a minimum at $|\tilde{\phi}|_{\rm min} \simeq \left(\frac{v^2}{g}\right)^{\frac{1}{n}}$ and $\text{Im}\ \tilde{\phi}^n_{\rm min} = 0$ with negative energy density $V(\tilde{\phi}_{\rm min}) \simeq -3\mathcal{A}(13)\exp\left(|\tilde{\phi}|^2\right)\left|\mathcal{W}_1(\tilde{\phi}_{\rm min})\right|^2 \simeq -3\mathcal{A}(13)\left(\frac{n}{n+1}\right)^2 v^4 \left|\tilde{\phi}_{\rm min}\right|^2$. However, in the context of SUGRA, we may interpret that such negative potential energy is almost canceled by positive contribution due to the local supersymmetry breaking, $\Lambda^4_{\rm SUGRA}$, and that the residual positive energy density is responsible for the present dark energy. Then, we can relate the energy scale of this model to the gravitino mass as $m_{3/2} \simeq \frac{n}{n+1}\left(\frac{v^2}{g}\right)^{\frac{1}{n}} v^2$. Identifying the real part of $\tilde{\phi}$ with the inflaton $\phi \equiv \sqrt{2}\text{Re}\tilde{\phi}$, the dynamics of the inflaton is governed by the following potential,

$$V^{(4)}(\phi) \simeq \mathcal{A}(13)\left(v^4 - \frac{2g}{2^{n/2}}v^2\phi^n + \frac{g^2}{2^n}\phi^{2n}\right). \tag{3.16}$$

Imposing renormalization condition, here we restrict ourselves to $n=2$ leading to effective $\mathcal{N}=1, \mathcal{D}=4$ SUGRA potential

$$V^{(4)}(\phi) = \mathcal{A}(13)\left(v^4 - gv^2\phi^2 + \frac{g^2}{4}\phi^4\right). \tag{3.17}$$

IV. MODELING INFLATION FROM DBI GALILEON

Summing up all the contributions from eqn(3.1), eqn(3.6), eqn(3.7) and eqn(3.13), the model for *D3 DBI Galileon* is described by the following effective action:

$$\begin{aligned} S^{(4)}_{Total} = \int d^4x \sqrt{-g^{(4)}} \Big[&\hat{\tilde{K}}(\phi,\tilde{X}) - \tilde{G}(\phi,\tilde{X})\Box^{(4)}\phi + \tilde{l}_1 R_{(4)} \\ &+ \tilde{l}_4\left(\mathcal{C}(1) R^{\alpha\beta\gamma\delta(4)} R^{(4)}_{\alpha\beta\gamma\delta} - 4\mathcal{I}(2) R^{\alpha\beta(4)} R^{(4)}_{\alpha\beta} + \mathcal{A}(6) R^2_{(4)}\right) + \tilde{l}_3\Big], \end{aligned} \tag{4.1}$$

where

$$\begin{aligned} &\hat{\tilde{K}}(\phi,\tilde{X}) = \tilde{K}(\tilde{\phi},\tilde{X}) - 2X\tilde{M}(T,T^\dagger) - \tilde{Z}(T,T^\dagger)V^{(4)}(\phi), \tilde{M}(T,T^\dagger) = \tfrac{M(T,T^\dagger)}{2\kappa^2_{(4)}}, \tilde{Z}(T,T^\dagger) = \tfrac{Z(T,T^\dagger)}{2\kappa^2_{(4)}}, \\ &\tilde{l}_1 = \left\{\tfrac{1}{2\kappa^2_{(4)}}\left[1 + \tfrac{\alpha_{(4)}}{R^2\beta^2}\left(24\mathcal{I}(2) - 24\mathcal{A}(9) - 16\mathcal{A}(10)\right)\right] - \tfrac{\alpha_{(4)}\mathcal{C}(2)}{\kappa^2_{(4)}R^2\beta^2}\right\}, \tilde{l}_4 = \tfrac{\alpha_{(4)}}{2\kappa^2_{(4)}}, \\ &\tilde{l}_3 = \tfrac{1}{2\kappa^2_{(4)}}\left[\tfrac{\alpha_{(4)}}{R^4\beta^4}\left(24\mathcal{C}(24) - 144\mathcal{I}(4) - 64\mathcal{A}(5) + 144\mathcal{A}(7) + 64\mathcal{A}(8) + 192\mathcal{A}(11)\right) - \tfrac{3M_5^3\beta b_0^6}{2\kappa^2_{(4)}M^2_{PL}R^5}\mathcal{I}(1)\right] \end{aligned}. \tag{4.2}$$

The collective effect of equation(3.8) and equation(3.17) gives the total *D3 DBI Galileon* potential as:

$$V(\phi) = \sum_{m=-2, m\neq -1}^{2} C_{2m}\phi^{2m}, \tag{4.3}$$

where $C_0 = \left(T_3\tilde{\nu}_0 + \beta R\mathcal{I}(2)\tilde{f}_0 + \tilde{Z}(T,T^{\dagger})\mathcal{A}(13)v^4\right)$, $C_{-4} = T_3\tilde{\nu}_4$, $C_2 = \left(\beta R\mathcal{I}(2)\tilde{f}_2 - gv^2\tilde{Z}(T,T^{\dagger})\mathcal{A}(13)\right)$, $C_4 = \left(\beta R\mathcal{I}(2)\tilde{f}_4 + \frac{\tilde{Z}(T,T^{\dagger})\mathcal{A}(13)g^2}{4}\right)$ are tree level constants. Now including the contribution from one-loop radiative correction, the renormalizable potential is as under

$$V(\phi) = \sum_{m=-2, m\neq -1}^{2} C_{2m}\left[1 + D_{2m}\ln\left(\frac{\phi}{M}\right)\right]\phi^{2m}, \tag{4.4}$$

where $D_0 = 0$. It is the Coleman Weinberg potential [37], provided the coupling constant satisfies the Gellmann-Low equation [38] in the context of Renormalization group. Here the first term in the eqn(4.4) physically represents the energy scale of inflation ($\sqrt[4]{C_0}$).



FIG. 1: Variation of one loop corrected potential($V(\phi)$) versus inflaton field (ϕ)

Figure (1) represents the inflaton potential for different values of C_{2m} and D_{2m}. From the observational constraints the best fit model is given by the range: $5.67\times 10^{-11}M_{PL}^4 < C_0 < 6\times 10^{-11}M_{PL}^4$, $1.01\times 10^{-16}M_{PL}^8 < C_{-4} < 2\times 10^{-16}M_{PL}^8$, $7.27\times 10^{-10}M_{PL}^2 < C_2 < 7.31\times 10^{-10}M_{PL}^2$, $2.01\times 10^{-14} < C_4 < 2.45\times 10^{-14}$, $0.014 < D_{-4} < 0.021$, $0.002 < D_2 < 0.012$ and $0.011 < D_4 < 0.019$ so that while doing numericals, we shall restrict ourselves to this range. Here $M_{PL} = 2.43\times 10^{18}GeV$ represents reduced 4D Planck mass. Consequently the energy scale of inflation has a window $0.658\times 10^{16}GeV < \sqrt[4]{C_0} < 0.667\times 10^{16}GeV$ which precisely falls within the GUT scale.

Now the action (4.1) leads to 4D effective *Einstein-Gauss Bonnet* equation as follows:

$$G^{(4)}_{\alpha\beta} + \alpha_{(4)}H^{(4)}_{\alpha\beta} = 8\pi G_{(4)}T^{(4)}_{\alpha\beta} - \Lambda_{(4)}g^{(4)}_{\alpha\beta} \tag{4.5}$$

where the *Energy Momentum tensor*, *Einstein tensor* and *Gauss-Bonnet tensor* are given by

$$\begin{aligned} T^{(4)}_{\alpha\beta} = T^{(bulk)(4)}_{\alpha\beta} + T^{(brane)(4)}_{\alpha\beta} = -2X - \tfrac{\mathcal{S}_{(1)}R}{b_0A_1}g^{(4)}\left\{-Xg^{\rho\sigma(4)} + \tfrac{e^5_{\dot{5}}V^{(4)}_{Can}}{16\pi^2R^4\beta^2}\right\} \\ + \hat{\tilde{K}}_X\nabla_\alpha\phi\nabla_\beta\phi + \tilde{K}g_{\alpha\beta} - 2\nabla_{(\alpha}\tilde{G}_{\beta)}\phi + g_{\alpha\beta}\nabla_\lambda\tilde{G}\nabla^\lambda\phi \\ - \tilde{G}_X\Box\phi\nabla_\alpha\phi\nabla_\beta\phi \end{aligned} \tag{4.6}$$

$$G^{(4)}_{\alpha\beta} = \tilde{h}_1R^{(4)}_{\alpha\beta} - \tfrac{1}{2}\tilde{h}_2g^{(4)}_{\alpha\beta}R_{(4)}g^{(4)}_{\alpha\beta}\left\{6\tilde{h}_4 + 4\tilde{h}_5 - 3\tilde{h}_3\right\} \tag{4.7}$$

$$\begin{aligned}
H^{(4)}_{\alpha\beta} = &\Big\{R^{(4)}_{\alpha\rho\gamma\delta}R^{(4)}_{\beta\lambda\sigma\xi}\tilde{h}_7 + \tfrac{\tilde{h}_8}{R^2\beta^2}\Big[R^{(4)}_{\beta\lambda\sigma\xi}\Big(g^{(4)}_{\gamma\rho}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\rho}\Big) + R^{(4)}_{\beta\lambda\sigma\xi}\Big(g^{(4)}_{\gamma\rho}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\rho}\Big)\Big] + \\
&\tfrac{1}{R^2\beta^2}\Big(g^{(4)}_{\gamma\rho}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\rho}\Big)\Big(g^{(4)}_{\gamma\rho}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\rho}\Big)\Big\}g^{\rho\lambda(4)}g^{\gamma\sigma(4)}g^{\delta\xi(4)} - \Big\{R^{(4)}_{\alpha\lambda\beta\sigma}R^{(4)}_{\sigma'\rho}\tilde{h}_{11} \\
&+ \tfrac{\tilde{h}_{12}}{R^2\beta^2}R^{(4)}_{\sigma'\rho}\Big(g^{(4)}_{\lambda\beta}g^{(4)}_{\alpha\sigma} - g^{(4)}_{\alpha\beta}g^{(4)}_{\sigma\lambda}\Big) - \tfrac{3\tilde{h}_{13}}{R^2\beta^2}g^{(4)}_{\sigma'\rho} - \tfrac{3\tilde{h}_{14}}{R^4\beta^4}g^{(4)}_{\sigma'\rho}\Big(g^{(4)}_{\lambda\beta}g^{(4)}_{\alpha\sigma} - g^{(4)}_{\alpha\beta}g^{(4)}_{\sigma\lambda}\Big)\Big\}g^{\lambda\sigma'(4)}g^{\sigma\rho(4)} \\
&- \Bigg\{R^{(4)}_{\alpha\gamma}R^{(4)}_{\beta\delta}\tilde{h}_{16} + \tfrac{9\tilde{h}_{17}}{\beta^4R^4}g^{(4)}_{\alpha\gamma}g^{(4)}_{\beta\delta} - \tfrac{3\left(\tilde{h}_{18}+\tilde{h}_{19}\right)}{R^2\beta^2}\Bigg\}g^{\gamma\delta(4)} + R_{(4)}R^{(4)}_{\alpha\beta}\tilde{h}_{21} - \tfrac{1}{R^2\beta^2}\Big\{3R_{(4)}g^{(4)}_{\alpha\beta}\tilde{h}_{22} \\
&+ 12R^{(4)}_{\alpha\beta}\Big(\tilde{h}_{23} + \tilde{h}_{25}\Big) + \tfrac{36\left(\tilde{h}_{24}+\tilde{\tilde{h}}_{25}\right)}{R^4\beta^4}g^{(4)}_{\alpha\beta}\Bigg\} - \tfrac{1}{2}g^{(4)}_{\alpha\beta}\Bigg[\Bigg\{R^{(4)}_{\rho\sigma\lambda\eta}R^{(4)}_{\alpha\beta\gamma\delta}\tilde{h}_{26} + \tfrac{R^{(4)}_{\alpha\beta\gamma\delta}}{R^2\beta^2}\tilde{h}_{27}\Big(g^{(4)}_{\lambda\sigma}g^{(4)}_{\rho\eta} - g^{(4)}_{\rho\lambda}g^{(4)}_{\eta\sigma}\Big) \\
&+ \tfrac{R^{(4)}_{\rho\sigma\lambda\eta}}{R^2\beta^2}\tilde{h}_{28}\Big(g^{(4)}_{\gamma\beta}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\beta}\Big) + \tfrac{\tilde{h}_{29}}{R^4\beta^4}\Big(g^{(4)}_{\lambda\sigma}g^{(4)}_{\rho\eta} - g^{(4)}_{\rho\lambda}g^{(4)}_{\eta\sigma}\Big)\Big(g^{(4)}_{\gamma\beta}g^{(4)}_{\alpha\delta} - g^{(4)}_{\alpha\gamma}g^{(4)}_{\delta\beta}\Big)\Bigg\}g^{\rho\alpha(4)}g^{\sigma\beta(4)}g^{\gamma\lambda(4)}g^{\delta\eta(4)} \\
&- 4g^{\alpha\sigma(4)}g^{\beta\rho(4)}\Big\{R^{(4)}_{\sigma\rho}R^{(4)}_{\alpha\beta}\tilde{h}_{31} - \tfrac{3}{R^2\beta^2}\Big(R^{(4)}_{\alpha\beta}g^{(4)}_{\sigma\rho}\tilde{h}_{32} + R^{(4)}_{\sigma\rho}g^{(4)}_{\alpha\beta}\tilde{h}_{33}\Big) + \tfrac{9}{R^4\beta^4}g^{(4)}_{\sigma\rho}g^{(4)}_{\alpha\beta}\Big\} + \tfrac{1}{R^4\beta^4}\Big(144\tilde{h}_{40} \\
&- 64\tilde{h}_{36} + 192\tilde{h}_{41} + 64\tilde{h}_{42}\Big) - \tfrac{1}{R^2\beta^2}R_{(4)}\Big(24\tilde{h}_{38} + 16\tilde{h}_{39}\Big) + \tilde{h}_{37}\left(R_{(4)}\right)^2\Bigg]
\end{aligned} \tag{4.8}$$

From FLRW metric in 4D, equation(4.5) results in the following *Friedmann equations*:

$$H^4\tilde{g}_1 + H^2\tilde{g}_2 + \dot{H}H^2\tilde{g}_3 + \dot{H}^2\tilde{g}_4 + \dot{H}\tilde{g}_5 = \tilde{\Lambda}_{(4)} + 8\pi G_{(4)}\rho \tag{4.9}$$

$$H^4\tilde{f}_1 + H^2\tilde{f}_2 + \dot{H}H^2\tilde{f}_3 + \dot{H}^2\tilde{f}_4 + \dot{H}\tilde{f}_5 = -\tilde{\tilde{\Lambda}}_{(4)} + 8\pi G_{(4)}p \tag{4.10}$$

where the constants have been listed in appendix. The *energy density* and *pressure* are now given by:

$$\rho = \Big[2\tilde{K}_X X - \tilde{K} + 6H\tilde{G}_X\dot{\phi}X - 2\tilde{G}_\phi X - 2X\left(1 - \Theta_1\right) + \Theta_3(T,T^\dagger)V_{bulk}(\phi)\Big], \tag{4.11}$$

$$p = \Big[\tilde{K} - 2\left(\tilde{G}_\phi + \tilde{G}_X\ddot{\phi}\right)X - 2X\Theta_1 - \Theta_3\left(T,T^\dagger\right)V_{bulk}(\phi)\Big], \tag{4.12}$$

where $\Theta_1 = \frac{\mathcal{S}_1 R}{b_0 A_1}$ and $\Theta_3(T,T^\dagger) = \frac{\sqrt{2}}{16\pi^2 b_0 R^3(T+T^\dagger)}$. Now in the slow-roll regime the *Galileon effects* can alter the dynamics. Hence *Friedmann* and *Klein-Gordon* equations can be expressed as:

$$H^4 = \frac{\Lambda_{(4)} + 8\pi G_{(4)}V(\phi)}{\tilde{g}_1}, \tag{4.13}$$

$$\dot{\phi}^2\left(e_2(\phi) + 9e_3(\phi)H^2\right) = \left\{V'(\phi) + \tilde{C}_5\tilde{g}'(\phi)k_1 - \frac{\tilde{D}\tilde{f}'(\phi)}{\tilde{f}(\phi)}\left(1 - Q_1\right)\right\}, \tag{4.14}$$

where $e_2(\phi) = \tilde{M}(T,T^\dagger)J_\phi + 2\tilde{g}(\phi)\tilde{f}'(\phi)k_1k_2 + 8\tilde{f}(\phi)\tilde{f}'(\phi)\tilde{g}(\phi)k_1k_2^2 + 2\tilde{g}'(\phi)\tilde{f}(\phi)k_1k_2 - g''(\phi)k_1$ and $e_3(\phi) = 2\tilde{C}_4\tilde{f}(\phi)\tilde{g}(\phi)k_1k_2$ provided $|e_3(\phi)| \gg |e_2(\phi)|$ in the slow-roll regime. Here we have fixed the signature of $\dot{\phi}$ so that the scalar field rolls down the potential. Additionally ghost instabilities are avoided provided the coefficient of $\dot{\phi}^2 > 0$. Consequently the potential dependent slow-roll parameters can be expressed as:

$$\epsilon_V := \frac{M_{PL}^2}{2}\left(\frac{V'}{V}\right)^2\frac{1}{\sqrt{\mathcal{G}(\phi)V'(\phi)}}, \tag{4.15}$$

$$\eta_V := M_{PL}^2\left(\frac{V''}{V}\right)\frac{1}{\sqrt{\mathcal{G}(\phi)V'(\phi)}}, \tag{4.16}$$

$$\xi_V := M_{PL}^4\left(\frac{V'V'''}{V^2}\right)\frac{1}{\mathcal{G}(\phi)V'(\phi)}, \tag{4.17}$$

$$\sigma_V := M_{PL}^6\left(\frac{(V')^2V''''}{V^3}\right)\frac{1}{\left(\mathcal{G}(\phi)V'(\phi)\right)^{\frac{3}{2}}}, \tag{4.18}$$

where $\mathcal{G}(\phi) = \frac{16e_3(\phi)M_{PL}^2}{\tilde{g}_1 V(\phi)}$. In this connection Galileon terms effectively flatten the potential due to the presence of the *flattening factor* $\frac{1}{\sqrt{\mathcal{G}(\phi)V'(\phi)}} \ll 1$. This implies that in the presence of Galileon like derivative interaction slow-roll inflation can take place even if the potential is rather steep.

The number of e-foldings for *D3 DBI Galileon* can be expressed as

$$\mathcal{N} = \frac{1}{8M_{PL}} \int_{\phi_i}^{\phi_f} \frac{\sqrt[4]{\mathcal{G}(\phi)V'(\phi)}}{\sqrt{\epsilon_V}} d\phi, \tag{4.19}$$

where ϕ_i and ϕ_f are the corresponding values of the inflaton field at the beginning and end of inflation.



FIG. 2: Variation of the number of e-foldings (N) vs inflaton field (ϕ)

Figure(2) represents a graphical behavior of number of e-folding versus the inflaton field for different values of $C_i^{'}s$ and the most satisfactory point in this context is the number of e-folding lies within the observational window $56 < N < 70$. The end of the inflation leads to the extra constraint $V^{'}(\phi_f) = \sqrt{V''(\phi_f)V(\phi_f)}$.

V. QUANTUM FLUCTUATIONS AND OBSERVABLE PARAMETERS

Let us now engage ourselves in analyzing quantum fluctuation in our model and its observational imprints via primordial spectra generated from cosmological perturbation. To serve this purpose, following the well known *ADM formalism*, the action can be expanded in the second order as:

$$S^{(4)}_{Total}|_S = \int dt d^3x a^3 Y_S \left[\dot{\zeta}^2 - \frac{c_s^2}{a^2}(\partial\zeta)^2\right], \tag{5.1}$$

where ζ comes from the ADM metric at the linear level and carries the signature of curvature perturbation. In equation(5.1)

$$Y_S = \frac{t_1\left(4t_1t_3 + 9t_2^2\right)}{3t_2^2}, \qquad c_s^2 = \frac{3\left(2Ht_2t_1^2 - t_4t_2^2 - 2t_1^2\dot{t}_2\right)}{t_1\left(4t_1t_3 + 9t_2^2\right)} \tag{5.2}$$

with $t_1 = \tilde{l}_1$, $t_2 = \left(2H\tilde{l}_1 - 2\dot{\phi}X\tilde{G}_X\right)$ and $t_3 = -9\tilde{l}_1H^2 + 3\left(X\hat{\tilde{K}}_X + 2X^2\hat{\tilde{K}}_{XX}\right) + 18H\dot{\phi}\left(2X\tilde{G}_X + X^2\tilde{G}_{XX}\right)$. It is important to mention here that *ghosts* and *Laplacian* instabilities can be avoided iff $c_s^2 > 0, Y_S > 0$. For convenience, we have introduced a new parameter defined as $\epsilon_s = \frac{Y_S c_s^2}{\tilde{l}_1}$.

Now varying the action stated in equation(5.1) and expressing the solution at the linear level in terms of Fourier modes, we arrive at the *Mukhanov Sasaki Equation* for Galileon scalar mode.

$$v_{\vec{k}}^{''} + \left(c_s^2k^2 - \frac{z^{''}}{z}\right)v_{\vec{k}} = 0, \tag{5.3}$$



(a)

(b)

(c)

FIG. 3: (a)Parametric plot of the amplitude of the power spectrum ($\sqrt{\mathcal{P}_\zeta}$) vs scalar spectral index (n_ζ), (b)Variation of the scale dependent scalar power spectrum ($\mathcal{P}_\zeta(k)$) vs momentum scale k, (c)Variation of the scale dependent tensor to scalar ratio ($r(k\eta)$) vs $|k\eta|$ for *DS* and *BDS* analysis.

where c_s^2 takes into account the nontrivial modification due to Galileon. Similarly for tensor modes, equation(5.1) can be recast as:

$$S^{(4)}_{Total}|_T = \int dtd^3xa^3Y_T\left[\dot{h}_{ij}^2 - \frac{c_s^2}{a^2}(\partial h_{ij})^2\right], \tag{5.4}$$

where

$$Y_T = \frac{t_1}{4} = \frac{\tilde{l}_1}{4}, \qquad c_T^2 = \frac{t_4}{t_1} = 1 + \mathcal{O}(\epsilon^2). \tag{5.5}$$

Following the same prescription we can establish equation(5.3) for tensor modes provided c_s is replaced by c_T. The *Bunch-Davies* mode function turns out to be (Throughout the paper we have used *DS* for *de-Sitter* results and *BDS* for *beyond de-Sitter* results.)

$$u(\eta,k) = \begin{cases} \dfrac{iH\exp(-ikc_s\eta)}{2\sqrt{Y_S}(c_sk)^{\frac{3}{2}}}(1+ikc_s\eta) & :DS \\ \dfrac{\sqrt{-k\eta c_s}}{a\sqrt{2Y_S}}\mathcal{H}^{(1)}_{\nu_s}(-k\eta c_s) & :BDS\,. \end{cases} \tag{5.6}$$

where $\nu_s = \left(\frac{3-\epsilon_V - 2s^S_V + \delta_V}{2(1-\epsilon_V - s^S_V)}\right)$ and in the *super-Hubble* limit $\mathcal{H}^{(1)}_{\nu_s} \rightarrow \frac{(-kc_s\eta)^{-\nu_s}\exp\left(i[\nu_s - \frac{1}{2}]\frac{\pi}{2}\right)2^{\nu_s - \frac{3}{2}}}{\sqrt{2c_s k}}\left(\frac{\Gamma(\nu_s)}{\Gamma(\frac{3}{2})}\right)$. Further we have introduced five new parameters during the analysis of primordial quantum fluctuation defined as:

$$\begin{aligned} s^S_V &:= \frac{\dot{c_s}}{Hc_s} = \frac{4\sqrt{V'(\phi)}M^2_{PL}}{\sqrt{\mathcal{G}(\phi)}}\frac{d}{d\phi}\left(\ln c_s\right), \quad (5.7)\\ s^T_V &:= \frac{\dot{c_T}}{Hc_T} = \frac{M_{PL}\sqrt{\tilde{g}_1 V'(\phi)}}{2\sqrt{e_3(\phi)V(\phi)}}\frac{d}{d\phi}\left(\ln\left[1+\mathcal{O}(\epsilon^2_V)\right]\right), \quad (5.8)\\ \eta_s &:= \frac{\dot{\epsilon_s}}{H\epsilon_s} = \frac{4\sqrt{V'(\phi)}M^2_{PL}}{\sqrt{\mathcal{G}(\phi)}}\frac{\left[\epsilon^{'}_V\left(1+\mathcal{O}(\epsilon_V)\right) + \delta_{GX\phi}\right]}{\left[\epsilon_V + \delta_{GX} + \mathcal{O}(\epsilon^2_V)\right]}, \quad (5.9)\\ \delta_V &:= \frac{\dot{Y_s}}{HY_s} = \frac{4\sqrt{V'(\phi)}M^2_{PL}}{\sqrt{\mathcal{G}(\phi)}}\frac{d}{d\phi}\left(\ln Y_s\right), \quad (5.10)\\ \delta_{GX} &:= \frac{\dot{\phi}X\tilde{G}_X}{\tilde{l}_1}. \quad (5.11)\end{aligned}$$

Now using eqn(5.6) the *two-point* correlation function for scalar modes can be expressed as:

$$\langle 0|\zeta(\vec{k})\zeta(\vec{k^{'}})|0\rangle = \frac{2\pi^2}{k^3}(2\pi)^3\mathcal{P}_\zeta(k)\delta^3(\vec{k}+\vec{k^{'}}), \quad (5.12)$$

where the *dimensionless Power spectrum* for scalar modes $\mathcal{P}_\zeta(k)$ at the horizon crossing turns out to be:

$$\mathcal{P}_\zeta(k_\star) = \begin{cases} \left(\frac{\sqrt{V(\phi)}}{8\pi^2 c_s\epsilon_s\tilde{l}_1\sqrt{\tilde{g}_1}M_{PL}}\right)_\star & \text{:DS} \\ \left(2^{2\nu_s-3}\left|\frac{\Gamma(\nu_s)}{\Gamma(\frac{3}{2})}\right|^2\frac{\left(1-\epsilon_V - s^S_V\right)^2\sqrt{V(\phi)}}{8\pi^2 Y_S c^3_s\sqrt{\tilde{g}_1}M_{PL}}\right)_\star & \text{:BDS .}\end{cases} \quad (5.13)$$

$\star$ corresponds to the horizon crossing. Similarly the *dimensionless Power spectrum* for tensor modes reads:

$$\mathcal{P}_T(k_\star) = \begin{cases} \left(\frac{\sqrt{V(\phi)}}{2\pi^2 c_T\epsilon_T\tilde{l}_1\sqrt{\tilde{g}_1}M_{PL}}\right)_\star & \text{:DS} \\ \left(2^{2\nu_T-3}\left|\frac{\Gamma(\nu_T)}{\Gamma(\frac{3}{2})}\right|^2\frac{\left(1-\epsilon_V - s^T_V\right)^2\sqrt{V(\phi)}}{2\pi^2 Y_T c^3_T\sqrt{\tilde{g}_1}M_{PL}}\right)_\star & \text{:BDS .}\end{cases} \quad (5.14)$$

Consequently the ratio of tensor to scalar power spectrum can be expressed as:

$$r(k_\star) = \frac{\mathcal{P}_T(k_\star)}{\mathcal{P}_\zeta(k_\star)} = \begin{cases} \left(16\epsilon_s c_s\left[1-\frac{3}{2}\mathcal{O}(\epsilon^2_T)\right]\right)_\star & \text{:DS} \\ \left(16.2^{2(\nu_T-\nu_s)}\left|\frac{\Gamma(\nu_T)}{\Gamma(\nu_s)}\right|^2\left(\frac{1-\epsilon_V - s^T_V}{1-\epsilon_V - s^S_V}\right)^2 c_s\epsilon_s\left[1-\frac{3}{2}\mathcal{O}(\epsilon^2_T)\right]\right)_\star & \text{:BDS .}\end{cases} \quad (5.15)$$

Further, the scale dependence of the perturbations, described by the scalar and tensor spectral indices, as follows:

$$n_\zeta - 1 = \left(\frac{d\ln\mathcal{P}_\zeta}{d\ln k}\right)_\star = \begin{cases} \left(-2\epsilon_V - \eta_s - s^S_V\right)_\star = \left(-2\epsilon_s - \eta_s - s^S_V + 2\delta_{GX} + 2\mathcal{O}(\epsilon^2_s)\right)_\star & \text{:DS} \\ (3-2\nu_s) = -\left(\frac{2\epsilon_V + s^S_V + \delta_V}{1-\epsilon_V - s^S_V}\right)_\star & \text{:BDS .}\end{cases} \quad (5.16)$$

$$n_T = \left(\frac{d\ln\mathcal{P}_T}{d\ln k}\right)_\star = \begin{cases} -2\epsilon_V|_\star = \left(-2\epsilon_s + 2\delta_{GX} + 2\mathcal{O}(\epsilon^2_s)\right)_\star & \text{:DS} \\ (3-2\nu_T) = -\left(\frac{s^T_V}{1-\epsilon_V - s^T_V}\right)_\star & \text{:BDS .}\end{cases} \quad (5.17)$$

Consistently, the consistency relation is also modified to:

$$r = \begin{cases} \left(8c_s\left(2\epsilon_V + 2\delta_{GX} + 2\mathcal{O}(\epsilon_V^2)\right)\left[1 - \frac{3}{2}\mathcal{O}(\epsilon_T^2)\right]\right)_\star & \textit{:DS} \\ \left(8.2^{2(\nu_T - \nu_s)}\left|\frac{\Gamma(\nu_T)}{\Gamma(\nu_s)}\right|^2\left(\frac{1-\epsilon_V - s_V^T}{1-\epsilon_V - s_V^S}\right)^2 c_s\left(2\epsilon_V + 2\delta_{GX} + 2\mathcal{O}(\epsilon_V^2)\right)\left[1 - \frac{3}{2}\mathcal{O}(\epsilon_T^2)\right]\right)_\star & \textit{:BDS}\,. \end{cases} \tag{5.18}$$

The expressions for the running of the scalar and tensor spectral index in this specific model with respect to the logarithmic pivot scale at the horizon crossing are given by:

$$\alpha_\zeta = \left(\frac{dn_\zeta}{d\ln k}\right)_\star = \begin{cases} \left\{\frac{4\sqrt{V'(\phi)}M_{PL}^2}{\sqrt{\mathcal{G}(\phi)}}\left(-2\epsilon_V^{'} - \eta_s^{'} - s_V^{S\,'}\right)\right\}_\star & \textit{:DS} \\ \frac{4\sqrt{V'(\phi)}M_{PL}^2}{\sqrt{\mathcal{G}(\phi)}\left(1-\epsilon_V - s_V^S\right)^2}\left[\underbrace{s_V^{S\,'}\epsilon_V^{'}} + \underbrace{\delta_V^{'}\epsilon_V^{'}} + \underbrace{s_V^{S\,'}\delta_V^{'}} + \left(2\epsilon_V^{'} + s_V^{S\,'} + \delta_V^{'}\right)\right] & \textit{:BDS}\,. \end{cases} \tag{5.19}$$

$$\alpha_T = \left(\frac{dn_T}{d\ln k}\right)_\star = \begin{cases} -2\left\{\frac{4\sqrt{V'(\phi)}M_{PL}^2}{\sqrt{\mathcal{G}(\phi)}}\epsilon_V^{'}\right\}_\star & \textit{:DS} \\ -\left[\frac{4\sqrt{V'(\phi)}M_{PL}^2}{\sqrt{\mathcal{G}(\phi)}}\left(\frac{s_V^{T\,'}}{1-\epsilon_V - s_V^T}\right) + \frac{s_T\left(\epsilon_V^{'} + s_V^{T\,'}\right)}{\left(1-\epsilon_V - s_V^T\right)^2}\right]_\star & \textit{:BDS}\,. \end{cases} \tag{5.20}$$

Here we have used a shorthand notation $\underbrace{ab} = a^{'}b - ab^{'}$ where $' = \frac{d}{d\phi}$.

Figure(3(a)) represents the scale dependent power spectrum ($\sqrt{\mathcal{P}_\zeta}$) with respect to the scalar spectral index(n_ζ). It directly shows that both the *DS* and *BDS* analysis follow the same characteristics but the estimated windows for the observational parameters ($\mathcal{P}_\zeta, n_\zeta$) are slightly different, but both of them are within the observational bound. Figure(3(b)) shows the characteristic differences between the behavior of *DS* and *BDS* scale dependent power spectrum with respect to the momentum scale (k). Here *DS* behavior is quasi-statically flat, but *BDS* characteristics is rapidly increasing with respect to the scale. In figure(3(c)) we have plotted the the scale dependent tensor to scalar ratio for *DS* and *BDS* limit. Most significantly they show complementary characteristics with the scale and intersects at a point where both the analysis will be equivalent. In the next section, we will estimate these parameters by confronting the results directly to WMAP7 results.

VI. PARAMETER ESTIMATION AND CONFRONTATION WITH WMAP7

Using the parameter space for the model parameters (C_i, D_i) we have estimated the window of the cosmological parameters from our model which confronts observational data well in $56 < N < 70$. In Table(I) we have tabulated the relevant observational parameters estimated from our model for both *DS* and *BDS* limit.

Scheme	$\sqrt{\mathcal{P}_\zeta}$ $\times 10^{-9}$	r	n_ζ	α_ζ $\times\left(-10^{-3}\right)$
DS	2.401 - 2.601	0.215 - 0.242	0.964 - 0.966	2.240 - 2.249
BDS	2.471 - 2.561	0.232 - 0.250	0.962 - 0.964	4.008 - 4.012

TABLE I: Model Dependent Observational Parameters

H_0 km/sec/MPc	τ_{Reion}	$\Omega_b h^2$	$\Omega_c h^2$	T_{CMB} K
71.0	0.09	0.0226	0.1119	2.725

TABLE II: Input parameters in CAMB

t_0 Gyr	z_{Reion}	Ω_m	Ω_Λ	Ω_k	η_{Rec} Mpc	η_0 Mpc
13.707	10.704	0.2670	0.7329	0.0	285.10	14345.1

TABLE III: Output parameters from CAMB

Further, we use the publicly available code CAMB [33] to verify our results directly with observation. To operate CAMB at the pivot scale $k_0 = 0.002Mpc^{-1}$ the values of the initial parameter space are taken for lower bound of $C_i^{'}$s and $N = 70$. Additionally WMAP7 years dataset for ΛCDM background has been used in CAMB to obtain CMB angular power spectrum. In Table(II) we have given all the input parameters for CAMB. Table(III) shows the CAMB output, which is in good agreement with WMAP7 [34] data. In figure(4)(a)-figure(4)(c) we have plotted CAMB output of CMB TT, TE and EE angular power spectrum $C_l^{TT}, C_l^{TE}, C_l^{EE}$ for the best fit with WMAP7 data for scalar mode, which explicitly show the agreement of our model with WMAP7 dataset. The small scale modes have no impact in the CMB anisotropy spectrum only the large scale modes have little contribution and this is obvious from figure(4)(d)-figure(4)(f) where we have plotted the CAMB output of CMB angular power spectrum C_l^{TT}, C_l^{TE}, C_l^{EE} and C_l^{BB} for best fit with WMAP7 data for the tensor mode. Hence in figure(5)we have plotted the variation of matter power spectrum with respect to the momentum scale which is in concordance with observational results.



(a) (b) (c)

(d) (e) (f)

FIG. 4: Variation of the CMB (a)TT (scalar), (b)TE (scalar), (c)EE (scalar), (d)EE+BB (tensor), (e)TT (tensor) and (f)TE (tensor) angular power spectrum with multipoles (l).

VII. SUMMARY AND OUTLOOK

In this article we have proposed a model of single field inflation in the context of DBI Galileon cosmology in D3 brane. We have demonstrated the technical details of construction mechanism of an one-loop 4D inflationary potential via dimensional reduction starting from D4 brane in $\mathcal{N}$=2,$\mathcal{D}$=5 SUGRA including the quadratic *Gauss-Bonnet* correction term that leads to an effective $\mathcal{N}$=1,$\mathcal{D}$=4 SUGRA in the D3 brane, which is precisely the DBI Galileon in our framework. Hence we have studied inflation using the one loop effective potential by estimating the observable parameters originated from primordial quantum fluctuation for scalar and tensor modes, in the de-Sitter



FIG. 5: Variation of matter power spectrum vs momentum scale k

and beyond de-sitter limit. We have further confronted our results with WMAP7 [34] dataset by using the cosmological code CAMB. The results are found in good agreement with WMAP7 dataset in $\Lambda CDM{+}tens$ background.

An interesting open issue in this context is to study the primordial non-Gaussian features of DBI Galileon introduced in the present article. As has been pointed out recently [39]-[41] there is a tension between bispectrum (f_{NL}) and tensor-to-scalar ratio (r) in DBI inflation, which is a generic sensitivity problem. It will be interesting to investigate whether our proposed framework of DBI Galileon can resolve this issue. We are already in progress in this direction and have obtained some interesting results. A detailed report on this issue will be brought forth shortly.

Other open issues in the context of DBI Galileon cosmology are combined constraints on the primordial non-Gaussianity via preheating [42], reheating [43] and primordial black hole formation [44], effect of the presence of one loop and two loop radiative corrections in the presence of all possible scalar and tensor mode fluctuations up to the fourth order correction in the action, study of different shapes and comparative study between the tree, one and two loop level via rigorous analysis and finding out the most probable Dark Matter candidate in collider phenomenology. It will be interesting to see how they are affected due to the presence of Galileon.

Acknowledgments

SC thanks Council of Scientific and Industrial Research, India for financial support through Junior Research Fellowship (Grant No. 09/093(0132)/2010).

Appendix

In this section we have explicitly mentioned the expressions for the constants appearing in the section III and IV.

$$\mathcal{I}(1)=\int_{-\pi R}^{+\pi R}dy\frac{4\left(3\exp(2\beta y)+3T_4^2\exp(-2\beta y)-2T_4\right)}{R^2\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^2}, \tag{7.1}$$

$$\mathcal{C}(1)=\frac{R^7}{b_0^7}\int_{-\pi R}^{+\pi R}dy\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{7}{2}}, \tag{7.2}$$

$$\mathcal{I}(2)=\mathcal{A}(6)=\tilde{h}_{11}=\frac{R^3}{b_0^3}\int_{-\pi R}^{+\pi R}dy\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{3}{2}}, \tag{7.3}$$

$$\mathcal{C}(2)=\frac{\beta^2R^5}{4b_0^5}\int_{-\pi R}^{+\pi R}dy\frac{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{5}{2}}\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^2}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^2}, \tag{7.4}$$

$$\mathcal{C}(4) = \mathcal{I}(4) = \mathcal{A}(5) = \frac{\beta^4 R^3}{16 b_0^3} \int_{-\pi R}^{+\pi R} dy \frac{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{3}{2}} \left(T_4 \exp(-\beta y) - \exp(\beta y)\right)^4}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^4}, \tag{7.5}$$

$$\mathcal{A}(7) = \frac{\beta^4 b_0^5}{16 R^5} \int_{-\pi R}^{+\pi R} dy \frac{\left(T_4 \exp(-\beta y) - \exp(\beta y)\right)^4}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^4 \left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}}, \tag{7.6}$$

$$\mathcal{A}(8) = \frac{4\beta^4 b_0^5 T_4^2}{R^5} \int_{-\pi R}^{+\pi R} dy \frac{1}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^4 \left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}}, \tag{7.7}$$

$$\mathcal{A}(9) = \tilde{h}_4 = \tilde{h}_{30} = \frac{\beta^2 b_0^5}{4R^5} \int_{-\pi R}^{+\pi R} dy \frac{\left(T_4 \exp(-\beta y) - \exp(\beta y)\right)^2}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^2 \left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}} \tag{7.8}$$

$$\mathcal{A}(10) = \tilde{h}_5 = -\frac{2\beta^2 b_0^5 T_4}{R^5} \int_{-\pi R}^{+\pi R} dy \frac{1}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^2 \left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}}, \tag{7.9}$$

$$\mathcal{A}(11) = -\frac{2\beta^4 b_0^5 T_4}{2R^5} \int_{-\pi R}^{+\pi R} dy \frac{\left(T_4 \exp(-\beta y) - \exp(\beta y)\right)^2}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^4 \left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}}, \tag{7.10}$$

$$\mathcal{A}(12) = -\frac{b_0}{R} \int_{-\pi R}^{+\pi R} dy \frac{\exp\left(-\frac{8\pi i y}{R}\right)}{\sqrt{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)}}, \tag{7.11}$$

$$\mathcal{A}(1) = \tilde{h}_1 = \frac{b_0}{R} \int_{-\pi R}^{+\pi R} dy \frac{1}{\sqrt{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)}}, \tag{7.12}$$

$$\mathcal{A}(13) = \frac{b_0 \tilde{h}_{16}}{R} = \frac{b_0 \tilde{h}_{31}}{R} = \int_{-\pi R}^{+\pi R} dy \sqrt{\exp(\beta y) + T_4 \exp(-\beta y)}, \tag{7.13}$$

$$\tilde{h}_2 = \frac{b_0^5}{R^5} \int_{-\pi R}^{+\pi R} dy \frac{1}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}}, \tag{7.14}$$

$$\tilde{h}_3 = \frac{b_0 \beta^2}{4R} \int_{-\pi R}^{+\pi R} dy \frac{\left(T_4 \exp(-\beta y) - \exp(-\beta y)\right)^2}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}}, \tag{7.15}$$

$$\tilde{h}_6 = \tilde{h}_{21} = \frac{b_0^3}{R^3} \int_{-\pi R}^{+\pi R} dy \frac{1}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{3}{2}}}, \tag{7.16}$$

$$\tilde{h}_7 = \tilde{h}_{26} = \frac{R^5}{b_0^5} \int_{-\pi R}^{+\pi R} dy \left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{5}{2}}, \tag{7.17}$$

$$\tilde{h}_8 = \tilde{h}_{13} = \tilde{h}_{27} = \tilde{h}_{28} = \frac{\beta^2 R^3}{4 b_0^3} \int_{-\pi R}^{+\pi R} dy \frac{\left(T_4 \exp(-\beta y) - \exp(\beta y)\right)^2}{\left(\exp(\beta y) + T_4 \exp(-\beta y)\right)^{\frac{1}{2}}}, \tag{7.18}$$

$$\tilde{h}_9=\tilde{h}_{14}=\tilde{h}_{17}=\tilde{h}_{29}=\tilde{h}_{34}=\tilde{h}_{36}=\frac{\beta^4 R}{16b_0}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^4}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{7}{2}}}, \tag{7.19}$$

$$\tilde{h}_{10}=\tilde{h}_{12}=\tilde{h}_{15}=\tilde{h}_{18}=\tilde{h}_{19}=\tilde{h}_{20}=\tilde{h}_{32}=\tilde{h}_{33}=\tilde{h}_{35}=\frac{\beta^2 R}{4b_0}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^2}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{3}{2}}}, \tag{7.20}$$

$$\tilde{h}_{22}=\tilde{h}_{23}=\frac{\beta^2 b_0^3}{4R^3}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^2}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{7}{2}}}, \tag{7.21}$$

$$\tilde{h}_{24}=\frac{\beta^4 b_0^3}{64R^3}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^4}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{11}{2}}}, \tag{7.22}$$

$$\tilde{h}_{25}=-\frac{2\beta^2 T_4 b_0^3}{R^3}\int_{-\pi R}^{+\pi R}dy\frac{1}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{7}{2}}}, \tag{7.23}$$

$$\tilde{\tilde{h}}_{25}=-\frac{\beta^4 T_4 b_0^3}{2R^3}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^2}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{11}{2}}}, \tag{7.24}$$

$$\tilde{h}_{37}=\frac{b_0^7}{R^7}\int_{-\pi R}^{+\pi R}dy\frac{1}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{7}{2}}}, \tag{7.25}$$

$$\tilde{h}_{38}=\frac{\beta^2 b_0^7}{4R^7}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^2}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{11}{2}}}, \tag{7.26}$$

$$\tilde{h}_{39}=-\frac{2\beta^2 T_4 b_0^7}{R^7}\int_{-\pi R}^{+\pi R}dy\frac{1}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{11}{2}}}, \tag{7.27}$$

$$\tilde{h}_{40}=\frac{\beta^4 b_0^7}{16R^7}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^4}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{15}{2}}}, \tag{7.28}$$

$$\tilde{h}_{41}=-\frac{\beta^4 T_4 b_0^7}{2R^7}\int_{-\pi R}^{+\pi R}dy\frac{\left(T_4\exp(-\beta y)-\exp(\beta y)\right)^2}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{15}{2}}}, \tag{7.29}$$

$$\tilde{h}_{42}=\frac{4\beta^4 T_4^2 b_0^7}{R^7}\int_{-\pi R}^{+\pi R}dy\frac{1}{\left(\exp(\beta y)+T_4\exp(-\beta y)\right)^{\frac{15}{2}}}, \tag{7.30}$$

$$
\begin{aligned}
&\tilde{g}_1 = \tilde{\alpha}_1 + \tilde{\alpha}_4 + \tilde{\alpha}_5 + a^2\left(\tilde{\alpha}_6 + \tilde{\alpha}_7\right), \quad \tilde{g}_2 = \left(\tilde{\alpha}_2 + \tilde{\alpha}_3\right), \quad \tilde{g}_3 = 2\tilde{\alpha}_1 + \tilde{\alpha}_5 + a^2\left(\tilde{\alpha}_6 + 2\tilde{\alpha}_7\right),\\
&\tilde{g}_4 = \tilde{\alpha}_1 + a^2\tilde{\alpha}_7, \quad \tilde{g}_5 = \tilde{\alpha}_2,\\
&\tilde{f}_1 = \tilde{\beta}_1 + \tilde{\beta}_5 + \tilde{\beta}_6, \quad \tilde{f}_2 = \tilde{\beta}_2 + \tilde{\beta}_3, \quad \tilde{f}_3 = 2\tilde{\beta}_1 + \tilde{\beta}_6, \quad \tilde{f}_4 = \tilde{\beta}_1, \quad \tilde{f}_5 = \tilde{\beta}_3,\\
&\tilde{\alpha}_1 = \alpha_{(4)}\left(-3\tilde{h}_7 - 9\tilde{h}_{16} - 18\tilde{h}_{21} + \tfrac{3}{2}\tilde{h}_{26} - 18\tilde{h}_{31} + 36\tilde{h}_{39}\right),\\
&\tilde{\alpha}_2 = 3\left(\tilde{h}_2 - \tilde{h}_1\right) + \tfrac{\alpha_{(4)}}{R^2\beta^2}\Big(12\tilde{h}_8 - 3\tilde{h}_{12} + 9\tilde{h}_{13} + 9\tilde{h}_{18} + 9\tilde{h}_{19} + 18\tilde{h}_{22} + 36\tilde{h}_{23}\\
&\qquad\qquad\qquad\qquad + 24\tilde{h}_{25} + 18\tilde{h}_{32} + 18\tilde{h}_{33} - 72\tilde{h}_{38} - 48\tilde{h}_{39}\Big),\\
&\tilde{\alpha}_3 = 3\tilde{h}_2 - \tfrac{\alpha_{(4)}}{R^2\beta^2}\left(18\tilde{h}_{22} - 6\tilde{h}_{12} - 72\tilde{h}_{38} - 48\tilde{h}_{39}\right), \qquad \tilde{\alpha}_4 = 36\alpha_{(4)}\tilde{h}_{39},\\
&\tilde{\alpha}_5 = \alpha_{(4)}\left(72\tilde{h}_{39} - 18\tilde{h}_{21}\right), \qquad \tilde{\alpha}_6 = 6\alpha_{(4)}\tilde{h}_{11}, \qquad \tilde{\alpha}_7 = 3\alpha_{(4)}\tilde{h}_{11},\\
&\tilde{\Lambda}_4 = \Lambda_4 - \tfrac{\alpha_{(4)}}{R^4\beta^4}\left(32\tilde{h}_{42} + 96\tilde{h}_{41} + 72\tilde{h}_{40} - 64\tilde{h}_{36} - 18\tilde{h}_{34} - 24\tilde{\tilde{h}}_{25} - 36\tilde{h}_{24} - 9\tilde{h}_{17} - 6\tilde{h}_9\right) - \tfrac{9\tilde{h}_{14}}{R^2\beta^2},\\
&\tilde{\beta}_1 = \alpha_{(4)}\left(\tilde{h}_7 - 3\tilde{h}_{11} - \tilde{h}_{16} + 6\tilde{h}_{21} - \tfrac{\tilde{h}_{26}}{2} + 2\tilde{h}_{31} - 3\tilde{h}_{37}\right),\\
&\tilde{\beta}_2 = \left(2\tilde{h}_1 - 3\tilde{h}_2\right) + \alpha_{(4)}\Big\{\tfrac{1}{R^2\beta^2}\Big(-8\tilde{h}_8 + 4\tilde{h}_{12} + 6\tilde{h}_{13} + 6\tilde{h}_{18} + 6\tilde{h}_{14} - 18\tilde{h}_{22} - 24\tilde{h}_{23}\\
&\qquad\qquad\qquad - 16\tilde{h}_{25} - 12\tilde{h}_{32} - 12\tilde{h}_{33} + 72\tilde{h}_{38} + 48\tilde{h}_{39}\Big) - \tilde{h}_{26}\Big\},\\
&\tilde{\beta}_3 = \left(\tilde{h}_1 - 3\tilde{h}_2\right) + \tfrac{\alpha_{(4)}}{R^2\beta^2}\Big(-4\tilde{h}_8 + 5\tilde{h}_{12} + 3\tilde{h}_{13} + 3\tilde{h}_{18} + 3\tilde{h}_{19} - 18\tilde{h}_{22} - 12\tilde{h}_{23}\\
&\qquad\qquad\qquad\qquad - 8\tilde{h}_{25} - 6\tilde{h}_{32} - 6\tilde{h}_{33} + 72\tilde{h}_{38} + 48\tilde{h}_{39}\Big),\\
&\tilde{\beta}_5 = \alpha_{(4)}\left(-3\tilde{h}_{37} + 8\tilde{h}_{31} + 12\tilde{h}_{21} - 4\tilde{h}_{16} - 4\tilde{h}_{11} + 2\tilde{h}_7\right),\\
&\tilde{\beta}_6 = \alpha_{(4)}\left(-3\tilde{h}_{37} + 8\tilde{h}_{31} + 18\tilde{h}_{21} - 4\tilde{h}_{16} - 2\tilde{h}_{11}\right),\\
&\tilde{\tilde{\Lambda}}_4 = \left(6\tilde{h}_4 + 4\tilde{h}_5 - 3\tilde{h}_3 + \Lambda_4\right) + \tfrac{\alpha_{(4)}}{R^4\beta^4}\Big(72\tilde{h}_{40} + 96\tilde{h}_{41} + 64\tilde{h}_{42} - 64\tilde{h}_{36} + 9\tilde{h}_{34}\\
&\qquad\qquad\qquad\qquad + 24\tilde{\tilde{h}}_{25} + 36\tilde{h}_{24} - 9\tilde{h}_{17} - 9\tilde{h}_{14} + 6\tilde{h}_9\Big).
\end{aligned}
\tag{7.31}
$$

$$
\begin{aligned}
&Q_1 = \frac{\mathcal{I}(3)}{\left(\bar{C}_1 d_2 - \bar{C}_2 d_4\right)}, \quad Q = \frac{\left(\bar{C}_2 d_3 - \bar{C}_1 d_1\right)}{\left(\bar{C}_1 d_2 - \bar{C}_2 d_4\right)},\\
&\bar{C}_1 = \frac{b_0\exp(-2\beta\pi R)}{2R\beta\left[\left(\frac{4\pi}{R}\right) + i\beta\right]\left[\left(\frac{4\pi}{R}\right) + 5i\beta\right]T_4^{\frac{3}{2}}}\sqrt{\frac{\exp(-2\beta\pi R) + T_4}{\exp(-\beta\pi R) + T_4\exp(\beta\pi R)}},\\
&\bar{C}_2 = \frac{b_0\exp(-4i\beta\pi^2)}{2R\beta\left[\left(\frac{4\pi}{R}\right) + i\beta\right]\left[\left(\frac{4\pi}{R}\right) + 5i\beta\right]T_4^{\frac{3}{2}}}\sqrt{\frac{\exp(2\beta\pi R) + T_4}{\exp(\beta\pi R) + T_4\exp(-\beta\pi R)}},\\
&\mathcal{I}(3) = \frac{b_0}{R}\int_{-\pi R}^{+\pi R} dy\,\frac{\exp\left(-\frac{2\pi i y}{R}\right)}{\sqrt{\left(\exp(\beta y) + T_4\exp(-\beta y)\right)}},\\
&d_2 = 6\beta\exp(2i\pi^2)\left[\left(\beta - \frac{4i\pi}{R}\right)\ {}_2F_1\left[\frac{3}{2};\frac{5}{4} - \frac{\pi i}{R\beta};\frac{9}{4} - \frac{\pi i}{R\beta};-\frac{\exp(-2\beta\pi R)}{T_{(4)}}\right]\right.\\
&\qquad\left. + \left(5\beta - \frac{4i\pi}{R}\right)T_4\exp(2\beta\pi R)\ {}_2F_1\left[\frac{3}{2};\frac{1}{4} - \frac{\pi i}{R\beta};\frac{5}{4} - \frac{\pi i}{R\beta};-\frac{\exp(-2\beta\pi R)}{T_{(4)}}\right]\right],\\
&d_4 = 6\beta\left[\left(\beta - \frac{4i\pi}{R}\right)\ {}_2F_1\left[\frac{3}{2};\frac{5}{4} - \frac{\pi i}{R\beta};\frac{9}{4} - \frac{\pi i}{R\beta};-\frac{\exp(2\beta\pi R)}{T_{(4)}}\right]\right.\\
&\qquad\left. + \left(5\beta - \frac{4i\pi}{R}\right)T_4\ {}_2F_1\left[\frac{3}{2};\frac{1}{4} - \frac{\pi i}{R\beta};\frac{5}{4} - \frac{\pi i}{R\beta};-\frac{\exp(2\beta\pi R)}{T_{(4)}}\right]\right],\\
&d_3 = \left\{\frac{16\pi^2}{R^2} + \frac{24\pi i\beta}{R} - 5\beta^2\right\}\exp\left[\left(\beta + \frac{2i\pi}{R}\right)\pi R\right]\ {}_2F_1\left[\frac{3}{4};\frac{3}{2};\frac{7}{4};-\frac{\exp(2\beta\pi R)}{T_{(4)}}\right],\\
&d_1 = \left\{\frac{16\pi^2}{R^2} + \frac{24\pi i\beta}{R} - 5\beta^2\right\}\exp(\beta\pi R)\ {}_2F_1\left[\frac{3}{4};\frac{3}{2};-\frac{7}{4};-\frac{\exp(-2\beta\pi R)}{T_{(4)}}\right],
\end{aligned}
\tag{7.32}
$$

---

[1] P. J. E. Peebles and B. Ratra, Rev. Mod. Phys. 75 (2003) 559.

[2] S. M. Caroll, Living Rev. Rel. 4 (2001) 1.

[3] G. R. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B, 485 (2000) 208.
[4] C. Deffayet, G. R. Dvali, and G. Gabadadze, Phys. Rev. D 65 (2002) 044023.
[5] N. Arkani Hamed, H. C. Cheng, M. A. Luty and S. Mukohyama, JHEP 0405 (2004) 074.
[6] G. Dvali, hep-th/0402130.
[7] A. I. Vainshtein, Phys. Lett. B, 39 (1972) 393.
[8] A. D. Felice and S. Tsujikawa, Living Rev. Rel. 13 (2010) 3.
[9] E. Babichev, C. Deffayet and R. Ziour, JHEP 0905 (2009) 098.
[10] G. F. R. Ellis, R. Maartens and M. A. H. MacCallum, Relativistic Cosmology, Cambridge University Press, 2012.
[11] C. Deffayet, G. E. Farese and A. Vikman, Phys. Rev. D 79 (2009) 084003.
[12] A. Nicolis, R. Rattazzi and E. Trincherini, Phys. Rev. D 79 (2009) 064036.
[13] A. D. Felice and S. Tsujikawa, Phys. Rev. Lett. 105 (2010) 111301.
[14] F. P. Silva and K. Koyama, Phys. Rev. D 80 (2009) 121301.
[15] N. Chow and J. Khoury, Phys. Rev. D 80 (2009) 024037.
[16] T. Kobayashi, M. Yamaguchi and J. Yokoyama, Phys. Rev. Lett. 105 (2010) 231302.
[17] M. Cederwall, A. von Gussich, A. Mikovi, b, B. E. W. Nilsson and A. Westerberg, Phys. Lett. B 390 (1997) 148.
[18] E. J. Copeland, S. Mizuno and M. Shaeri, Phys. Rev. D 81 (2010) 123501.
[19] C. de Rahm and A. J. Tolley, JCAP 1005 (2010) 015.
[20] G. L. Goon, K. Hinterbichler and M. Trodden, Phys. Rev. D 83 (2011) 085015.
[21] G. L. Goon, K. Hinterbichler and M. Trodden, JCAP 07 (2011) 017.
[22] A. De Felice and S. Tsujikawa, JCAP 02 (2012) 007.
[23] A. De Felice, K. Karwan and P. Wongjun, Phys. Rev. D 85 (2012) 123545.
[24] M. Ostrogradski, Mem. Ac. St. Petersbourg VI 4, 385 (1850).
[25] E. Babichev, C. Deffayet and R. Ziour, Int. J. Mod. Phys. D 18 (2009) 2147.
[26] E. Babichev, C. Deffayet and G. E. Farese, Phys. Rev. D 84 (2011) 061502.
[27] T. Kobayashi, M. Yamaguchi and J. Yokoyama, Prog. Theor. Phys. 126 (2011) 511.
[28] K. Kamada, T. Kobayashi, M. Yamaguchi and J. Yokoyama, Phys. Rev. D 83 (2011) 083515.
[29] C. Deffayet, O. Pujolas, I. Sawicki and A. Vikman, JCAP 1010 (2010) 026.
[30] S. Choudhury and S. Pal, Phys. Rev. D 85 (2012) 043529.
[31] S. Choudhury and S. Pal, Nucl. Phys. B 857 (2012) 85.
[32] H. P. Nilles, Phys. Rept. (1984) 110.
[33] CAMB, Online link: *http://camb.info/*.
[34] WMAP collaboration, D. N. Spergel et al., Astrophys. J. Suppl. **170**, 377 (2007); for uptodate results on WMAP, see *http://lambda.gsfc.nasa.gov/product/map/current*.
[35] Planck collaboration, *http://www.rssd.esa.int/index.php?project=Planck* , some early results are also available, see, for example, P. A. R. Ade et.al., arXiv:1101.2022.
[36] M. Yamaguchi, arXiv:1101.2488.
[37] S. Coleman and E. Weinberg, Phys. Rev. D 7 (1973) 1888.
[38] L. H. Ryder, Quantum Field Theory, Cambridge University Press (2008).
[39] S. E. Shandera and S. H. Henry Tye, JCAP 0605 (2006) 007.
[40] S. H. Henry Tye, Lect. Notes Phys. 737 (2008) 949.
[41] R. Bean, S. E. Shandera, S. H. Henry Tye and J. Xu, JCAP 075 (2007) 004.
[42] D. K. Hazra, J. Martin and L. Sriramkumar, arXiv:1206.0442.
[43] T. Suyama and M. Yamaguchi, Phys. Rev. D 77 (2008) 023505.
[44] C. T. Byrnes, E. J. Copeland and A. M. Green, arXiv:1206.4188.